

The Fellows' Pelvic Research Network (FPRN)

Lowenstein L, Olivera C, Harvie H, Schaffer J, Young S
Int Urogynecol J Pelvic Floor Dysfunct. 2008 Jul;19(7):895-7.

Anatomic outcomes of vaginal mesh procedure (Prolift, Gynecare) compared to uterosacral ligament suspension and abdominal sacrocolpopexy for pelvic organ prolapse.

Sanses TVD., Shahryarinejad A., Molden S., Hoskey KA., Abbasy S., Patterson D., Saks EK., Weber-Lebrun E., Gamble, TL., King VG., Nguyen AL., Abed H., Young SB.
American Journal of Obstetrics & Gynecology. 2009 Nov;201(5):519.e1-8.

Sexual Function is Related to Body Image Perception in Women with Pelvic Organ Prolapse.

Lowenstein L, Gamble T, Deniseiko Sanses TV, van Raalte H, Carberry C, Jakus S, Kambiss S, McAchran S, Pham T, Aschkenazi S, Hoskey K, Kenton K.
Journal of Sexual Medicine. 2009 Aug; 6(8):2286-91

Sexual Activity and Function in Women with and without PFDs

T Omotosho, H Harvie, M Schimpf, L Epstein, C Olivera, M Jean-Michel, K Rooney, S Balgabon, R Gala, R Rogers. Submitted: American Journal of Obstetrics & Gynecology

Does Vaginal Size Impact Sexual Activity and Function?

M Schimpf, H Harvie, T Omotosho, L Epstein, M Jean-Michel, C Olivera, K Rooney, S Balgabon, R Gala, R Rogers. Int.Urogynecol J Pelvic Floor Dysfunct. 2010 Apr;21(4):447-52.

Changes in Sexual Function after Treatment for Prolapse Are Related to the Improvement in Body Image Perception.

Lowenstein L, Gamble T, Sanses TV, van Raalte H, Carberry C, Jakus S, Pham T, Nguyen A, Hoskey K, Kenton K. J Sex Med. 2010 Feb;7(2 Pt 2):1023-8.